

Event anisotropy of identified π^0 , photon and electron compared to charged π , K , p and deuteron in $\sqrt{s_{NN}} = 200$ GeV Au+Au at PHENIX

Masashi Kaneta[†] for the PHENIX Collaboration[‡]

[†] RIKEN-BNL Research Center, Brookhaven National Laboratory, Upton NY 11973-5000, USA

E-mail: kaneta@bnl.gov

Abstract. We report the recent results of event anisotropy analysis focused on v_2 in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions at PHENIX.

1. Introduction

Event anisotropy analysis is a powerful tool for studying properties of the early stage in high-energy heavy ion collisions. Recent results of event anisotropy analysis suggest that finite v_2 (second harmonic coefficient of Fourier expansion of azimuthal distributions) is present up to relatively high transverse momentum ($p_T \sim 3\text{--}4$ GeV/ c) [1]. Measurement of π^0 's can reach very high p_T and thus should permit study of the effect of hard processes, e.g. jet quenching, on v_2 . The dominant source of photons seen in the experiment is from meson decays (mainly from π^0 's), and there have been several measurements on v_2 of charged hadron and identified π^0 up to high p_T close to 10 GeV/ c . As a next step in the systematic study of v_2 at high p_T , we have investigated subtracting the hadronic decay contribution from the v_2 of the measured photons to determine the feasibility of seeing the direct or thermal photon v_2 . We have also investigated the measured electron spectrum, which includes contributions from hadron decays and conversion photons (referred to as “*photonic electrons*”). These effects are well studied [2, 3] and we extract the charmed electron distribution after subtractions. We adopt the same method in electron v_2 analysis to obtain the charmed electron v_2 .

2. Analysis

Event anisotropy analysis is an application of the Fourier expansion of azimuthal particle distributions. Here we focus on the elliptic component of the event anisotropy. Due to momentum conservation and rotational symmetry, the odd order harmonics will be

[‡] For the full PHENIX Collaboration author list and acknowledgments, see Appendix “Collaboration” of this volume.

canceled when we add positive and negative rapidity regions. Additionally assuming that the contribution of higher order of harmonics is negligible, the distributions are described by

$$E \frac{dN^3}{d^3p} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left(1 + 2v_2^m \cos[2(\phi - \Phi_r)] \right) = C \left(1 + 2v_2^m \cos[2(\phi - \Phi_r)] \right), \quad (1)$$

where y is the rapidity and ϕ is the azimuthal angle of the particle, Φ_r is the reaction plane, v_2^m is the 2nd order of the Fourier coefficient measured by detectors, and C is a constant. The reaction plane is defined using Beam-Beam counters [4] ($|\eta|=3.1-4.0$) with a resolution due to signal size of v_2 and multiplicity on the detector. Therefore, v_2^m is broadened by the uncertainty of the reaction plane definition. The method of v_2^m correction we adopted, described in reference [5], results in a real v_2 corrected by the reaction plane resolution $v_2 = v_2^m / \sigma_2$, where σ_2 is the resolution for the 2nd harmonic. Since the reaction plane is distributed uniformly, the efficiency of the detector is canceled. Hence, we fit the number of particles as a function of $\phi - \Phi_r$ by equation (1) to obtain v_2^m .

The π^0 v_2 analysis is performed in the following steps: 1) counting the number of π^0 's from invariant mass from two photons measured by the EMCal [6, 7] as a function of $\phi - \Phi_r$ for each centrality and p_T bin, 2) fitting the distribution by equation (1), and 3) reaction plane resolution correction to v_2^m . The photon and electron v_2 analysis are checked by comparison to another method, which is $v_2^m = \langle \cos[2(\phi - \Phi_r)] \rangle$. The difference between two methods is taken into account as a systematic error. The additional contributions to the systematic error are the following: (1) Particle identification cut, (2) reaction plane determination, and (3) error propagation in “*photonic electron*” subtraction (only for charmed electron v_2). The total systematic error is estimated by adding the contributions quadratically.

3. Results and Discussion

Figure 1 shows photon and π^0 v_2 as a function of p_T from three centrality bins (top 20%, 20-40%, and 40-60%) in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions. The upper limit of the p_T range is defined by the available statistics while the lower limit of $p_T > 1$ GeV/ c results from hadron contamination in the EMCal. The π^0 v_2 increases with p_T and then saturates at $p_T \sim 3$ GeV/ c in each centrality. Comparison to the PHENIX data for identified charged pion v_2 [8] indicates good consistency with the π^0 v_2 results presented here (shown in reference [9]). Within errors, the photon v_2 is consistent with that of the π^0 v_2 . Better statistics is necessary to obtain the direct photon v_2 , especially at high p_T .

The result for π^0 v_2 shown in Figure 2 is the first measurement of an identified hadron up to $p_T = 10$ GeV/ c in minimum bias 200 GeV Au+Au collisions. It has a non-zero value up to $p_T = 8$ GeV/ c . The comparison is done with other hadrons from results of PHENIX [1, 10] and the STAR collaboration [11]. There is a crossover point among hadrons around $p_T = 2$ GeV/ c . The lower p_T (< 2 GeV/ c) region is well




Figure 1. (Color online) Preliminary results for photon (solid square) and π^0 (open diamond) v_2 as a function of p_T from three centrality bins in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions. The statistical error is shown as a vertical bar on each point. Systematic errors are shown by curves.




Figure 2. (Color online) Preliminary results for π^0 v_2 as a function of p_T from minimum bias data in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions, compared to other hadrons. The statistical error is shown as a vertical bar on each point and the systematic error is shown by curves for the π^0 's. The other hadrons are shown with statistical error (a vertical bar) only. The charged π , K , and p v_2 are from reference [1]. The deuteron v_2 is PHENIX preliminary data shown in reference [10]. The K_S^0 and Λ v_2 are from STAR in reference [11].

described by hydrodynamical models [1, 11]. In the region of $p_T > 2$ GeV/c, the pion v_2 merges with that of the kaon, as does the proton and Λ v_2 . At the same time, the meson v_2 remains different from that for baryons. This suggests a scaling rule called quark coalescence [12], related to the different number of quarks in mesons and baryons, which might indicate that the hadron v_2 is generated during the early partonic stage before the hadronization. A simple check of a coalescence picture in v_2 is normalization of both v_2 and p_T by the number of constituent quarks.

The left-hand side plot in Figure 3 is the plot to check the coalescence picture in v_2 . The data from π , K , p , Λ are superimposed and show good agreement in $p_T/n_{quark} > 1$ GeV/c. The plot shows that the deuteron v_2 agrees with the other hadrons by scaling number of quarks. However, we note that nucleon coalescence could be the dominant factor for forming deuterons rather than the quark coalescence; there is no way to separate them from deuteron v_2 measurements alone.




Figure 3. (Color online) The left-hand side plot shows v_2 as a function of p_T scaled via the coalescence prescription. The meson, Baryon, and deuteron are scaled by 2, 3, and 6, respectively. “Non-photonic” electron v_2 as a function of p_T is shown at the right-hand side. The vertical bar on each point show the statistical error and systematic errors are shown by colored box. The horizontal bar shows the RMS of dN/dp_T in each bin.

The “non-photonic” electron v_2 vs. p_T in minimum bias $\sqrt{s_{NN}} = 200$ GeV collisions is shown on the right-hand side of Figure 3 with two scenarios from a model calculation from reference [13]. One scenario is that D mesons are made from completely thermalized charm and light quarks (shown by solid line), while the other is that there is no interaction of the c -quarks, so that the flow contribution in D ’s is only from the light quarks (dashed line). The non-photonic electrons decay from charmed and bottomed mesons; those in the p_T range we showed are dominantly charmed meson decay. (A detailed discussion of the electron measurement is found in reference [3].) Within the available statistics, both models are consistent with data. It will be very interesting to pursue this comparison of charm flow with that of mesons containing light quarks using the higher statistics Run-4 data set.

References

- [1] PHENIX Collaboration K. Adcox *et al* 2003 *Preprint* nucl-ex/0307010
- [2] PHENIX Collaboration K. Adcox *et al* 2002 *Phys. Rev. Lett.* **88** 192303
- [3] PHENIX Collaboration S. Kelly *et al* in this proceedings
- [4] K. Ikematsu *et al* 1998 Nucl. Instrum. Meth. A **411** 238
- [5] A.M. Poskanzer and S.A. Voloshin 1998 *Phys. Rev. C* **58** 1671
- [6] PHENIX Collaboration S.S. Adler *et al* 2003 *Preprint* hep-ex/0304038
- [7] PHENIX Collaboration S.S. Adler *et al* 2003 *Phys. Rev. Lett.* **91** 072301
- [8] PHENIX Collaboration S.S. Adler *et al* 2003 *Preprint* nucl-ex/0305013
- [9] PHENIX Collaboration M. Kaneta *et al* 2003 DNP2003 talk.
- [10] PHENIX Collaboration S. Sakai *et al* 2003 HIS03 poster.
- [11] STAR Collaboration J. Adams *et al* 2004 *Phys. Rev. Lett.* **92** 052302
- [12] For example, V. Greco, C.M. Ko and P. Levai 2003 *Phys. Rev. C* **68** 034904 : R.J. Fries, B. Muller, C. Nonaka and S.A. Bass 2003 *Phys. Rev. Lett.* **90** 202303 : D. Molnár and S.A. Voloshin 2003 *Phys. Rev. Lett.* **91** 092301
- [13] V. Greco, C.M. Ko and R. Rapp 2003 *Preprint* nucl-th/0312100